

Vaccination Schedule 2020

More than 10 YEARS of Excellence

Disease	Vaccine	Birth	6 weeks	10 weeks	14 weeks	6 months	9 months	12 months	15 months	18 months	6 years	9 years	12 years
TB	BCG	●											
Polio	bOPV	●	●										
Diphtheria, Tetanus, Polio, Pertussis Haemophilus Influenzae, Hepatitis B	Hexaxim		●	●	●					●			
	Infanrix Hexa		●	●	●					●			
Pneumococcal	Prevenar 13		●		●								
	Synflorix		●	●	●		●	●	Nine and a half months				
Rotavirus	Rotarix		●		●								
	RotaTeq		●	●	●								
Measles	Measbio					●		●	12 months Measbio not required if giving MMR				
Measles Mumps Rubella	Priorix							●	MMR at 12 months instead of Measbio		●		
	OMZYTA							●	MMR at 12 months instead of Measbio		●		
Chickenpox	Varilrix								●		●		
	ONVARA								●		●		
Measles Mumps Rubella + Chickenpox	Priorix Tetra	Priorix Tetra can be used at 12 months instead of Measbio, MMR and Varicella. Increased risk of febrile convulsion after the first dose of Priorix Tetra							●		●		
Meningococcal Conjugate	Menactra						●		●	For individuals 2 - 55 years administer 1 dose			
Hepatitis A	AVAXIM 80							●		●			
	Havrix Junior							●		●			
Tetanus, Diphtheria Pertussis, Polio	TETRAxIM								Tetraxim at 6 years or Boostrix Tetra at 6 years		●		
	Boostrix Tetra								Boostrix Tetra can be given at 12 years if not given at 6 years. Adacel Quadra at 12 years		●		●
	ADACEL QUADRA								Adacel Quadra at 12 years		●		●
Tetanus, Diphtheria	Td										●		●
Human Papilloma Virus	Cervarix	Cervarix from 9 -14 years or Gardasil from 9 -13 years as 2 doses 6 months apart										●	
	GARDASIL	State programme: Cervarix to Grade 4 girls in public schools 9 -14 years. 2 doses 6 months apart										●	
Influenza	Influvac	From 6 months, important for children in creche, chronic illness and respiratory problems. Start before the INFLUENZA season in MAY.											
Respiratory Syncytial Virus	VAXIGRIp	From 6 months, important for children in creche, chronic illness and respiratory problems. Start before the INFLUENZA season in MAY.											
	SYNAGIS PALIVIZUMAB	RSV prophylaxis in high-risk infants-prevention of serious LRTI caused by RSV. Start January and end in May.											
Pneumococcal	PNEUMOVAX 23	Must have at least one dose of Conjugate PCV before Pneumovax 23. Pneumovax only in children older than 2 years with Immune Compromise or high risk of Pneumococcal infection. Two doses with 2nd dose 3 - 5 years after first.											

● STATE EPI VACCINES These vaccines are available free from Government supplied clinics

● RECOMMENDED OPTIONAL VACCINES Some of the vaccines in this schedule are only available from private clinics